

Review

Self-harm in older adults:
systematic review

M. Isabela Troya, Opeyemi Babatunde, Kay Polidano, Bernadette Bartlam, Erin McCloskey, Lisa Dikomitis and Carolyn A. Chew-Graham

Background

Self-harm is a major public health concern. Increasing ageing populations and high risk of suicide in later life highlight the importance of identification of the particular characteristics of self-harm in older adults.

Aim

To systematically review characteristics of self-harm in older adults.

Methods

A comprehensive search for primary studies on self-harm in older adults was conducted in e-databases (AgeLine, CINAHL, PsycINFO, MEDLINE, Web of Science) from their inception to February 2018. Using predefined criteria, articles were independently screened and assessed for methodological quality. Data were synthesised following a narrative approach. A patient advisory group advised on the design, conduct and interpretation of findings.

Results

A total of 40 articles ($n = 62\,755$ older adults) were included. Yearly self-harm rates were 19 to 65 per 100 000 people. Self-poisoning was the most commonly reported method. Comorbid physical problems were common. Increased risk repetition was

reported among older adults with self-harm history and previous and current psychiatric treatment. Loss of control, increased loneliness and perceived burdensome ageing were reported self-harm motivations.

Conclusions

Self-harm in older adults has distinct characteristics that should be explored to improve management and care. Although risk of further self-harm and suicide is high in all age cohorts, risk of suicide is higher in older adults. Given the frequent contact with health services, an opportunity exists for detection and prevention of self-harm and suicide in this population. These results are limited to research in hospital-based settings and community-based studies are needed to fully understand self-harm among older adults.

Declaration of interest

None.

Keywords

Self-harm; suicide; systematic review.

Copyright and usage

© The Royal College of Psychiatrists 2019.

Self-harm is a major public health concern worldwide, affecting not only those who self-harm but also family members and broader society through increased resource costs and productivity losses.^{1–3} In this review, self-harm is defined by National Institute for Health and Care Excellence (NICE) guidelines (CG16 and 133), as ‘any act of self-poisoning or self-injury carried out by a person, irrespective of motivation’.⁴ This review does not include indirect self-harm (e.g. refusal to eat/drink, self-neglect), but rather focuses on direct self-harm as defined by NICE guidelines (CG16 and 133).⁴ Self-harm and suicide are often linked to mental health problems; although self-harm and suicide can be seen as two distinct behaviours, self-harm is the major risk factor for suicide.^{5,6} The world’s population is ageing, and it is projected that 20% of the UK’s population will be 65 years and older by 2020.⁷ Rates of mental health conditions in later life are high (approximately 15% for adults aged 60 and over), and suicide rates are among the highest in older adults.^{8,9} An understanding of the nature of self-harm in later life is essential to offer more effective and adequate healthcare provision to this population. Previous reviews in the area were conducted over a decade ago, had no clear eligibility criteria for included studies and lacked quality appraisal of included studies.^{10,11} Consequently, this systematic review aimed to provide an up-to-date and robust synthesis of the evidence by describing the characteristics (rates and risk factors) of older adults who self-harm, including clinical characteristics and lived experiences of self-harm.

Methods

This review was conducted and reported in accordance with established systematic review guidance (Preferred Reporting Items for Systematic Reviews and Meta-Analyses, PRISMA). An *a priori* protocol was established and registered on PROSPERO, an international prospective register of systematic reviews (CRD42017057505).

Patient and Public Involvement and Engagement

The review was conducted in consultation with a Patient and Public Involvement and Engagement (PPIE) group, including members of a local self-harm group. A previous PPIE group had been convened for a former study on self-harm in primary care¹² and some members of the group had noted the importance of considering self-harm in older adults, resulting in the present study being conducted. Members of the original group expressing an interest in and experience of self-harm in older adults were reconvened. With over a decade of experience involving patients and the public in health research,¹³ this study was supported by the PPIE team at the Research Institute for Primary Care and Health Sciences at Keele University. All PPIE members were aged 60 or older, and included older adults with self-harm history, carers and support workers. The PPIE group was consulted four times at different stages of the review, including refining the review question, specification of study eligibility criteria, outcomes, interpretation and dissemination

of findings. The group also contributed to developing the diagrammatic representation of the relationship between the various risk factors for self-harm among older people (see *Results* section: *Influencing factors for self-harm*; Fig. 1). Findings based on lived experiences and current literature were discussed to reach consensus during PPIE meetings. These discussions were then considered when interpreting results from the review. Inclusion of the PPIE group was considered essential to ensure the study outcomes were mapped pragmatically to patient-centred outcomes.

Information sources, study selection and review process

A comprehensive search strategy was developed and used to search electronic databases (AgeLine, CINAHL, PsycINFO, MEDLINE and Web of Science) for published studies on self-harm in older adults. Databases were searched from their inception until 28 February 2018 (for full search strategy, see Supplementary Appendix 1 available at <https://doi.org/10.1192/bjp.2019.11>). Additionally, hand searching of reference lists of included studies was carried out to identify other potentially relevant grey literature. No language restrictions were applied.

Each identified study was evaluated against the following predetermined selection criteria.

- Population: Studies examining older adult populations (aged 60 years or older) with presence of at least one self-harm episode as defined by NICE.⁴
- Exposure: Self-harm determined by clinical presentation; self-report; or reports from family, carers or health practitioners regardless of suicidal or non-suicidal intent.
- Outcomes: Studies reporting at least one clinical characteristic (e.g. self-harm rates, methods, repetitions) and/or lived experiences (defined as an individual's representation and understanding of a particular experience¹⁴) with self-harm were included. Secondary outcomes such as specific diagnoses, mental illness and comorbidities, and personal demographics such as marital status and living conditions were highlighted but were not required for inclusion in the review.
- Study designs and settings: Observational studies with or without comparison groups from both clinical and community populations were included in the review.

Exclusion criteria were narrative reviews, letters, editorials, commentaries and conference abstracts for which there are no data and data requests were not successful. Case reports/case series and non-English language studies for which interpretation could not be obtained were also excluded.

The study selection process was tested and piloted *a priori* by members of the review team (M.I.T., K.P., B.B., O.B., C.A.C.-G.). Subsequently, two reviewers (M.I.T., K.P.) independently evaluated the eligibility of all identified citations. At each stage of title, abstract and full-text selection, disagreements regarding eligibility were resolved through discussion between reviewers (M.I.T., K.P.) or by the independent vote of a third reviewer (B.B., O.B. or C.A.C.-G.).

Data were extracted by one reviewer (M.I.T.) using a pre-tested customised data extraction form. Data were independently checked for completion, accuracy and consistency by a second reviewer (K.P. or E.M.). Data were extracted on the clinical characteristics of self-harm and lived experiences of the study participants. More specifically, data were extracted regarding population characteristics (e.g. age, gender, marital status, living situation, ethnicity), characteristics of self-harm including methods and rates, and outcomes (e.g. risk factors, clinical characteristics, contact with health services, motivations, stressors for self-harm). In instances of missing or incomplete quantitative data (i.e. lack of crude estimates or




Fig. 1 Influencing factors in self-harm in older adults.

Diagram presented in layers according to internal and external factors. Different size layers do not refer to higher or lower association to self-harm but rather represent internal and external factors.

measures of variability for estimates of self-harm), additional information was requested through contacting primary study authors. A random effects meta-analysis of quantitative self-harm data was planned but could not be performed due to inherent heterogeneity, incomplete reporting of data from primary studies and non-response to provision of required information from study authors. A descriptive analysis of quantitative data alongside a thematic analysis of qualitative data was performed and narratively synthesised together.¹⁵ Thematic analysis¹⁶ involved line-by-line coding, organisation of codes into descriptive themes and generation of analytical themes. Thematic analysis was conducted by one reviewer (M.I.T.) and then checked for completion, accuracy and consistency of identified themes by a second reviewer (E.M.).

Summary of evidence per risk factors for self-harm repetition were completed. A modified version of the Grading of Recommendations Assessment, Development and Evaluation (GRADE) rating system (<http://www.gradeworkinggroup.org/>) was used to assess the overall quality of evidence. The following factors were considered: the strength of association for each risk factor, methodological quality/design of the studies, consistency, directedness, precision, size and (where possible) dose-response gradient of the estimates of effects across the evidence base. Evidence was graded as very low, low, moderate and high, similar to a GRADE rating system.

The methodological quality of included studies was independently appraised by pairs of reviewers (M.I.T. and K.P. or O.B.), using the National Institutes of Health quality assessment toolkits for quantitative studies¹⁷ and the Critical Appraisal Skills Programme checklist for qualitative studies.¹⁸ Ratings of high, moderate or poor were given to studies according to the criteria stated in the toolkits. Disagreements regarding methodological quality of the included studies were resolved through discussion until consensus was reached.

Results

A total of 15 647 unique citations were identified, with 8 additional studies included through reference checking. A total of 405 abstracts were screened and 56 full-text articles were assessed for inclusion.




Fig. 2 Study flow diagram.

Forty studies (21 cross-sectional designs, 14 cohort studies, 3 qualitative studies and 2 case-control studies) met full eligibility criteria and were included. The flow of studies through the review process and reasons for exclusion are presented in Fig. 2.

Description of studies

Study setting

Country of origin of the included studies were mainly English-speaking countries ($n = 21$).^{19–39} However, 17 studies^{40–56} were from non-English-speaking countries, with 16 different countries being represented. Two were multi-site studies across Europe, including both English- and non-English-speaking countries.^{57,58} The majority of included studies were conducted in hospital-based settings ($n = 34$), mostly in emergency or psychiatry departments, with the exception of a plastic surgery department¹⁹ and a poisons unit.³⁴ The remaining studies were conducted in other healthcare facilities (e.g. general hospitals, general practice, private clinics) ($n = 2$),^{57,58} community mental health services ($n = 2$),^{32,50} a national surveillance system which includes presentations from hospitals and primary care⁴⁷ and a national household survey.⁵⁶ Study length varied from 8 months to 26 years. Follow-up was reported in all 14 cohort studies and varied from 1 to 23 years. All but one study⁵⁶ were based on self-harm presentations as determined by clinical presentation. The remaining study⁵⁶ was based on self-reported self-harm. The main characteristics of the included studies are summarised in Table 1.

Methodological quality assessment

Included studies were mostly of moderate ($n = 28$) to high ($n = 10$) methodological quality. Two studies were assessed as having poor quality. Figure 3a provides an overview of the quality assessment of studies and Fig. 3b highlights areas with higher or lower risk assessment. Risk assessment of studies was determined by grouping and rating the different methodological quality assessments of studies (e.g. confounding, loss to follow-up). High-risk ratings were given to studies where the quality assessment element was not reported at par with standards; studies were rated low risk when this was reported according to standards. Overall, participation rate, study population, research question, repeated exposure, time frame, defined outcomes and inclusion criteria were consistently assessed as having lower risk assessment across studies ($\geq 80\%$). Loss to follow-up and measurement and adjustment of confounding variables were rated as having higher risk across studies ($\geq 60\%$). Blinding of assessors and estimate of sample size were rated as having an unclear risk assessment across studies ($\geq 60\%$).

Self-harm outcomes

Sociodemographic characteristics

All but three studies^{30,38,55} ($n = 17\,377$) reported participants' gender: over half (57%; $n = 9903$) were women, and 43% ($n = 7474$) were men. Age of participants ranged from 60 to 112 years. Nine studies^{24,29,35,36,40,47,48,56,57} made a classification of

individuals according to age range ($n = 51\,174$): 60% ($n = 31\,072$) of participants were aged 60–74 years old. Eleven studies^{22,28,29,31,32,40,47,53–56} classified participants according to ethnicity ($n = 6573$), with the majority of participants being White (68.1%, $n = 4479$) and 13.3% ($n = 875$) were of other ethnicities (Black, Asian, Hispanic or Maori). The ethnicity of the remaining 18.6% ($n = 1219$) of participants was unknown. A total of 27 studies^{20–22,24,25,27,28,31–33,35,37,40,42–46,48–51,53–55,57,58} reported the marital status of their participants ($n = 4161$): approximately half were not married (51%, $n = 2121$), 38% ($n = 1582$) were married and the marital status of the remaining 11% ($n = 461$) was unknown. Over half of the studies^{19,20,23–28,31,35,41,44–46,48,50,51,53,55–58} ($n = 3103$) reported participants' living situation: 53.5% ($n = 1658$) were either living with family or in care, followed by 40% ($n = 1241$) who were living alone at the time of the self-harm event. The remaining of participants' living situation was unknown (6.5%, $n = 203$).

Self-harm rates

Overall, there were 63 266 self-harm presentations involving 62 755 older adult participants. Of the 40 included studies, 7^{23,25,27–29,36,57} presented overall estimates of self-harm rates per population ($n = 13\,776$). Yearly rates per 100 000 habitants varied from 19.3²⁹ to 65²³ as shown in Table 2.

Self-harm methods

Of the 40 included studies, 34 ($n = 61\,395$) reported self-harm methods used by older adults. Table 1 includes a summary of the reported methods, with the majority of self-harm presentations being self-poisoning (86.1%, $n = 52\,866$) which included overdose of medication or ingestion of toxic substances. Self-injury through lacerations or burning of skin was 8.1% ($n = 5002$). Other methods included hanging, gunshots, car fumes, jumping in front of cars and immolation (5.6%, $n = 3417$). The remaining 0.2% ($n = 110$) of participants used multiple methods to self-harm. The majority of studies reporting self-harm methods were in hospital-based settings, with the exception of four studies^{32,47,57,58} that also reported community-based data. However, similar trends regarding self-harm methods used were reported across the different study settings as reported in Table 1.

Associated clinical characteristics

Previous history of self-harm

A total of 30 studies^{19–28,31,32,34,35,37,39,40,42–46,48–50,53–55,57,58} reported previous history of self-harm ($n = 6033$). Nearly one third of participants (29.4%; $n = 1774$) had a previous history of self-harm.

Previous psychiatric history

A total of 30 studies^{19–29,31,32,35,37,39–46,48,51,54–58} reported participants' previous psychiatric history ($n = 10\,976$), including alcohol and substance misuse, schizophrenia and personality disorder, with 30% of participants having previous psychiatric history ($n = 3279$). Depression was the most commonly reported psychiatric diagnosis ($n = 7893$) across the 29 studies reporting depression. Specifically, 68.5% ($n = 5414$) of older adults who self-harmed had a diagnosis of depression.

Physical illness

A total of 25 studies^{20–28,31,34,35,37,39–42,45,48,49,51,53–55,58} reported comorbid physical illness among older adults who self-harm ($n = 4211$). Chronic physical illness (including cardiovascular disease,

diabetes, musculoskeletal disorders, neurological problems) was common among participants, with 40% having a comorbid condition ($n = 1666$).

Medication

Seven studies^{20,22,27,28,31,43,49} reported medication use by participants ($n = 689$). Nearly half of the participants from these studies (42.4%; $n = 292$) were prescribed antidepressants at the time of the self-harm episode.

Alcohol use

A total of 11 studies^{20,21,23,24,27,29,31,33,34,36,47} reported alcohol use at the time of the self-harm episode ($n = 13\,326$): 16% ($n = 2131$) of participants presenting with self-harm had consumed alcohol at the time of the episode.

Self-harm repetition and completed suicide

A total of 14 studies^{21–28,31,37,40,49,52,58} reported self-harm repetition ($n = 3065$). The time measurement period varied vastly from 1 to 23 years, and 17% ($n = 518$) of the older adult population that self-harmed repeated this behaviour during the study period.

The 16 studies^{21,23–28,31,33,37,39–41,49,52,58} that reported death of participants following self-harm ($n = 3883$) reflected this variation in follow-up time: up to 17% ($n = 653$) had died during the time of the studies. Not all of these studies specified causes of death, but in those that did ($n = 2939$), 3.3% ($n = 98$) died by suicide.^{21,23–26,31,37,40,41,49,52,58} As summarised in Table 1, the studies reporting self-harm repetition and completed suicide were all based in hospital settings.

Contact with health services

Contact with different health services ranging from primary care to specialised care such as psychiatric services were reported among participants in some of the studies.

Primary care

Three studies^{20,22,45} ($n = 208$) reported participants' previous contact with primary care services before self-harm episodes: 28.9% ($n = 42$) had seen their general practitioner 1 week before self-harming, whereas 62% ($n = 98$) had been in contact with primary care at least 1 month before the self-harm episode.

Psychiatric services

A total of 29 studies^{19–29,31,32,35,37,39–46,48,49,51,54,57,58} reported previous use of psychiatric services ($n = 5054$): 41.3% ($n = 2086$) of participants had previously attended services and/or received treatment before the self-harm episode. In contrast, only seven studies^{20–23,28,31,41} ($n = 2493$) reported participants receiving psychiatric treatment at the moment of the episode (28.2%; $n = 703$).

Follow-up

A total of 23 studies^{19–23,25–28,31,33–36,40,41,44,45,49,51,53–55} ($n = 8398$) reported that 52.4% ($n = 4403$) of participants received a psychiatric assessment immediately after the self-harm episode. Across the studies, there was no further follow-up or indication of whether this assessment led to any treatment or prevention of repeated self-harm.

Risk factors for self-harm repetition

Of the 40 included studies, 9^{21,23,31,35,37,49,53,55–56} calculated risk factors for self-harm repetition ($n = 2646$). The risk factors for

Table 1 Characteristics of included studies

Study ID	Study design	Quality assessment	Study setting	Study length (follow-up)	Participants (presentations)	Age range	Self-harm method	Self-harm repetition (time)	Death (suicide)	Influencing factors for self-harm	Motivations for self-harm
Armond 2017 ⁴⁷ Brazil	Cross-sectional	Moderate	Hospital and community based	1 year	93 (93)	60–90+	Self-poison: 39 Self-injury: 0 Other: 54	NA	NA	Low education level and socioeconomic status	NR
Bonnewyn 2014 ⁵¹ Belgium	Qualitative	Moderate	Hospital based (psychiatry)	NR	8 (8)	66–85	NR	NA	NA	Loss, death of spouse or family member, conflict with family member, physical illness, physical disability, insomnia, loneliness, loss of control	Death of a spouse or family member, conflict with family member, physical illness/disability, loneliness, loss of control
Briskman 2017 ⁴⁴ Israel	Cross-sectional	Moderate	Hospital based (A&E)	8 years	187 (187)	65–95	Self-poison: 177 Self-injury: 10 Other: 0	NA	NA	NR	NR
Carter 2014 ³⁶ USA	Cross-sectional	High	Hospital based (A&E)	1 year	4915 (4915)	65–85+	Self-poison: 3077 Self-injury: 595 Other: 1243	NA	NA	Alcohol and drug use	NR
Cheung 2017 ³¹ New Zealand	Cohort	High	Hospital based (A&E)	3 years (1 year)	339 (339)	65–96	Self-poison: 233 Self-injury: 31 Other: 37 Multiple: 38	50/339 (1 year)	7 (7)	Perceived physical illness, family discord, bereavement, financial trouble, partner separation	NR
Chiu 1996 ⁴⁰ Hong Kong	Cohort	Poor	Hospital based (psychiatry)	2.5 years (1.5 years)	55 (55)	65–91	Self-poison: 15 Self-injury: 40	5/55 (1.5 years)	16 (3)	NR	NR
Crocker 2006 ³² UK	Qualitative	High	Community based	NR	15 (15)	65–91	Self-poison: 14 Self-injury: 0 Other: 1	NA	NA	Social isolation, loss of social status, physical illness and loss of mobility, loneliness, ageing perceived as burdensome	Become invisible to others, regaining control
De Beer 2015 ²⁸ New Zealand	Cohort	Moderate	Hospital based (A&E)	3 years (1 year)	52 (52)	65–80+	Self-poison: 34 Self-injury: 8 Other: 7 Multiple: 3	7/52 (1 year)	5/52 (0)	Physical illness, pain, family discord, changed relationship, bereavement, financial trouble, legal difficulties	NR
De Leo 2001 ⁵⁷ Europe	Cross-sectional	Moderate	Hospital and community based	5 years	1518 (1734)	65–82	Self-poison: 1196 Self-injury: 191 Other: 347	NA	NA	NR	NR
De Leo 2002 ⁵⁸ Europe	Cross-sectional	Moderate	Hospital & community based	3 years (1 year)	63 (63)	60 and over ^a	Self-poison: 50 Self-injury: 3 Other: 4 Multiple: 6	15/63 (1 year)	8 (8)	Bereavement of father, poor mental health, poor social assistance, financial problems	Relational difficulties, desire to manifest desperation to others, mental illness
Dennis 2007 ²⁰ UK	Cross-sectional	Moderate	Hospital based (psychiatry)	NR	76 (76)	65–92	Self-poison: 43 Self-injury: 9 Other: 4 Multiple: 20	NA	NA	Isolated lifestyle, life events and difficulties, bereavement, health problems	Gain relief from an unbearable state of mind escape, make others understand how desperate they were, influence others, seek help, make others feel sorry
Draper 1994 ³⁵ Australia	Cross-sectional	Moderate	Hospital based (psychiatry)	6.5 years	69 (69)	65–85+	Self-poison: 52 Self-injury: 13 Multiple: 4	NA	NA	Social isolation, family issues, marital issues, death, accommodation issues, financial problems	NR
Gavrielatos 2006 ⁴³ Greece	Cross-sectional	Moderate	Hospital based (A&E)	3.5 years	44 (44)	65–91	Self-poison: 44 Self-injury: 0 Other: 0	NR	NR	Domestic stress (e.g. health or financial issues), stress of chronic illness	NR

(Continued)

Table 1 (Continued)

Study ID	Study design	Quality assessment	Study setting	Study length (follow-up)	Participants (presentations)	Age range	Self-harm method	Self-harm repetition (time)	Death (suicide)	Influencing factors for self-harm	Motivations for self-harm
Gheshlaghi 2012 ⁴¹ Iran	Cross-sectional	Poor	Hospital based (A&E)	1 year	43 (43)	65–83	Self-poison: 43 Self-injury: 0 Other: 0	NA	3 (3)	NR	NR
Gokcelli 2017 ⁴⁸ Turkey	Cross-sectional	Moderate	Hospital based (A&E)	9 years	63 (63)	60–91	Self-poison: 56 Self-injury: 3 Other: 4	NA	NA	NR	NR
Hawton 2006 ²¹ UK	Cohort	Moderate	Hospital based (psychiatry and A&E)	23 years (23 years)	730 (730)	60–85+	Self-poison: 647 Self-injury: 62 Multiple: 21	112/730 (23 years)	432 (30)	Physical illness, social isolation, relationship problems, bereavement, housing problems, alcohol misuse, financial worries	NR
Hepple 1997 ³⁷ UK	Cohort	Moderate	Hospital based (Psychiatry)	3 years (2–5 years)	100 (100)	65–94	Self-poison: 87 Self-injury: 2 Other: 11	28/100 (2–5 years)	42 (7)	Isolation, friction with family, bereavement, physical and psychiatric problems	NR
Kim 2011 ⁴² Korea	Cross-sectional	High	Hospital based (A&E)	2 years	57 (57)	65–81	Self-poison: 57 Self-injury: 0 Other: 0	NA	NA	NR	Interpersonal conflict, economic problems, physical illness
Kim 2014 ⁵⁰ Korea	Qualitative	Moderate	Community based	8 months	35 (35)	64–89	NR	NA	NA	Financial problems, domestic violence, illness, childhood events, violence, grief, mental illness	Feelings of helplessness, despair, dependence, isolation
Lamprecht 2005 ²² UK	Cohort	High	Hospital based (psychiatry)	3 years (1–2 years)	82 (99)	65–82	Self-poison: 90 Self-injury: 5 Other: 4	15/82 (1–2 years)	NR	Pain and debilitating illness	NR
Lawrence 2000 ³⁸ Australia	Cohort	Moderate	Hospital based	15 years	1368 (1596)	60–80+	NR	NA	NA	NR	NR
Lebret 2006 ⁴⁹ France	Cohort	High	Hospital based (psychiatry)	7 years (3 years)	59 (59)	61–85	Self-poison: 31 Self-injury: 9 Other: 12 Multiple: 7	8/59 (3 years)	17 (3)	Physical illness, loneliness, relationship conflict	Physical illness, interpersonal problems, social isolation/loneliness
Liu 2009 ⁵⁵ Taiwan	Case control	Moderate	Hospital based (A&E)	20 months	43 (43)	61–90	Self-poison: 21 Self-injury: 12 Other: 10	NA	NA	Health conditions, finances, interpersonal relations, affinity relations, parent–child relations	Depression, family conflict, long-term physical illness, financial burden
Logan 2007 ²⁹ USA	Cross-sectional	Moderate	Hospital based (A&E)	1 year	5710 (5710)	65 and over ^a	Self-poison: 3425 Self-injury: 1062 Other: 1223	NA	NA	NR	NR
Murphy 2012 ²³ UK	Cohort	High	Hospital based (A&E)	8 years (1–8 years)	1177 (1177)	60–97	Self-poison: 1031 Self-injury: 107 Other: 39	196/1177 (1 year)	24 (24)	Relationship problems, bereavement, physical and/or mental health problems, alcohol problems	NR
Nowers 1993 ²⁴ UK	Cohort	High	Hospital based (A&E)	7 years (5 years)	88 (88)	65–90	Self-poison: 85 Self-injury: 2 Other: 1	17/88 (1 year)	26 (5)	NR	NR
Packer 2012 ¹⁹ UK	Cross-sectional	Moderate	Hospital based (plastic surgery)	5 years	10 (10)	60–80+	Self-poison: 0 Self-injury: 10 Other: 0	NA	NA	NR	NR
Pierce 1987 ²⁵ UK	Cohort	Moderate	Hospital based (A&E)	13 years (1–12 years)	145 (145)	65–92	Self-poison: 138 Self-injury: 7 Other: 0	12/145 (1–12 years)	4 (4)	Physical illness, housing or financial stress, pain	NR

(Continued)

Table 1 (Continued)

Study ID	Study design	Quality assessment	Study setting	Study length (follow-up)	Participants (presentations)	Age range	Self-harm method	Self-harm repetition (time)	Death (suicide)	Influencing factors for self-harm	Motivations for self-harm
Pierce 1996 ²⁶ UK	Cohort	Moderate	Hospital based (A&E)	20 years (1–19 years)	39 (89)	60–87	Self-poison: 78 Self-injury: 6 Other: 5	39/39 (50 months)	18 (2)	NR	NR
Pillans 2017 ³⁹ Australia	Cross-sectional	High	Hospital based	26 years	626 (626)	65–97	Self-poison: 500 Self-injury: 126	NA	24 (NR)	NR	NR
Ruths 2005 ²⁷ UK	Cohort	Moderate	Hospital based (A&E)	2 years (2 years)	43 (43)	65–95	Self-poison: 36 Self-injury: 7 Other: 0	8/43 (2 years)	18 (0)	Chronic pain, terminal illness	NR
Shah 2009 ³⁰ UK	Cross-sectional	Moderate	Hospital based	9 years	44310 (44 310)	60–75+	Self-poison: 41 298 Self-injury: 2635 Other: 377	NA	NA	NR	NR
Takahashi 1995 ⁴⁵ Japan	Cross-sectional	Moderate	Hospital based (psychiatry)	10 years	50 (50)	65–89	Self-poison: 21 Self-injury: 13 Other: 5 Multiple: 11	NA	NA	NR	NR
Ticehurst 2002 ³³ Australia	Cross-sectional	Moderate	Hospital based	7.5 years	110 (110)	65 and over ^a	Self-poison: 110 Self-injury: 0 Other: 0	NA	6 (NR)	NR	NR
Tsoh 2005 ⁵³ Hong Kong	Case-control	Moderate	Hospital based (Psychiatry)	15 months	66 (66)	65–82	NR	NA	NA	Psychiatric and physical morbidities, family discord	NR
Van Orden 2015 ⁵² Sweden	Cohort	Moderate	Hospital based (A&E)	3 years (12 months)	101 (101)	70–91	Self-poison: 73 Self-injury: 13 Other: 15	6 (12 months)	2 (2)	Social problems, perceived to be burdensome, psychological problems, physical problems	Escape, functioning and autonomy, psychological problems, somatic problems and pain, perceived burden, social problems, lack of meaning
Wiktorsson 2010 ⁴⁶ Sweden	Cross-sectional	Moderate	Hospital based (A&E)	3 years	103 (103)	70–91	NR	NA	NA	Hopelessness, loneliness, low education	NR
Wynne 1987 ³⁴ UK	Cross-sectional	Moderate	Hospital based (Poisons unit)	4.5 years	45 (45)	65 and over ^a	Self-poison: 45 Self-injury: 0 Other: 0	NA	NA	Physical: pain, severe illness, disability, terminal illness Social: relationship, housing, financial problems Psychiatric: depression, personality disorder, alcohol problems	Physical: pain, severe illness, disability, terminal illness Social: relationship problems, housing, financial problems Psychiatric: depression, personality disorder, alcohol problems
Yang 2001 ⁵⁴ Taiwan	Cross-sectional	Moderate	Hospital based (psychiatry)	6 years	55 (55)	65–84	Self-poison: 20 Self-injury: 21 Other: 14	NA	NA	Psychosocial problems, family problems, interpersonal problems, adjustment problems, physical illness	Psychosocial problems, family problems, interpersonal problems, adjustment problems, physical illness
Zhang 2016 ⁵⁶ China	Cross-sectional	High	Community based	6 months	63 (63)	60–112	NR	NA	NA	Having no caregivers, psychological problems	NR

NA, not applicable; NR, not reported; A&E, accident and emergency.
a. Paper did not report upper range limit.

Table 2 Yearly self-harm in older adults rates per 100 000 habitants

Study	Study setting	Population size	Yearly rates per 100 000 habitants	CIs
Logan <i>et al</i> , 2007 ²⁹	Hospital based (A&E) US	<i>n</i> = 5710	19.3	95% CI 13.9–24.8
De Beer <i>et al</i> , 2015 ²⁸	Hospital based (A&E) New Zealand	<i>n</i> = 52	32.7 ^a	Not provided
Pierce 1987 ²⁵	Hospital based (A&E) UK	<i>n</i> = 145	46 ^a	Not provided
Ruths <i>et al</i> , 2005 ²⁷	Hospital based (A&E) UK	<i>n</i> = 43	47.3 ^a	Not provided
De Leo <i>et al</i> , 2001 ⁵⁷	Hospital and community based Multi-site study conducted in 13 countries in Europe	<i>n</i> = 1734	61.43	Not provided
Carter <i>et al</i> , 2014 ³⁶	Hospital based (A&E) US	<i>n</i> = 4915	63	95% CI 61.2–64.8
Murphy <i>et al</i> , 2012 ²³	Hospital based (A&E) UK	<i>n</i> = 1177	65	Not provided

A&E, accident and emergency.
a. Small population size (*n* < 200).

self-harm repetition (summarised below) are grouped according to sociodemographic, clinical or other factors. Table 3 provides a summary of findings per group for the identified risk factors for self-harm repetition.

Sociodemographic factors

Three studies estimated female gender to be a risk factor for self-harm repetition.^{21,35,49} Not being married or partnered, living alone and a younger age (being 60–74 years old) were also found to be risk factors.²³ Additionally, not having a caregiver was also found to be a risk factor for self-harm repetition.⁵⁶

Clinical factors

Previous episode of self-harm was found to be a risk factor for self-harm repetition among older adults.^{23,53} Three studies^{35,37,53} found that those with previous psychiatric history were also more likely to repeat self-harm. Four studies^{37,53,55,56} estimated that people with a depression diagnosis were more likely to repeat self-harm. In this review, both previous and current psychiatric treatment was found to be a risk factor for self-harm repetition in three studies.^{23,31,37} Finally, Tsoh and collaborators⁵³ also identified a diagnosis of arthritis as a risk factor for self-harm.

Other factors

Time was also found to be a determinant of self-harm repetition. Hawton and Harriss²¹ found that older adults were most likely to repeat self-harm within 12 months of the first episode. Two studies found alcohol and drug use as a risk factor for self-harm repetition.^{23,31} Poorer function of self-care was also found to be a risk factor for self-harm repetition.^{53,56}

Suicidal intention

Nine studies^{20–22,24,28,31,32,35,37} (*n* = 972) reported suicidal intention, with a total of 73.5% (*n* = 714) of participants declaring suicidal intent. A variety of tools to assess suicidal intention were used, including interviewer's assessment, questionnaires such as the Beck suicidal intent score and the Columbia Classification Algorithm of Suicide Assessment.

Motivations for self-harm

A total of 11 studies^{20,32,34,42,49–51,52,54,55,58} (*n* = 551; less than 1% of participants) presented motivations for self-harm with broader explanations besides suicidal intent. The identified motivations

emerged from both qualitative and quantitative studies and were based on self-reported motivations. Table 1 provides further detail of the identified motivations for self-harm which included relationship problems, physical and psychiatric illness, financial worries, regaining control, bereavement, isolation and helplessness.

Qualitative findings

Three qualitative studies^{32,50,51} (*n* = 58) explored lived experiences of self-harm in older adults. Participants had similar sociodemographic characteristics. Country of origin and study settings were diverse, including psychiatric departments,⁵¹ local mental health services^{32,50} and community groups for older adults.⁵⁰ The focus of the qualitative studies was self-harm with suicidal intention exclusively, as all studies classified the act of self-harm as a suicide attempt. Three major themes were identified consequent to data analysis: loss of control contributing to the suicide attempt, increased loneliness and isolation, and ageing perceived as burdensome and affecting daily living. Table 4 illustrates the three major themes with direct quotes of participants from the included articles.

Loss of control contributing to the suicide attempt was a major theme mentioned in two studies.^{32,51} Loss of control due to both physical and mental health problems was described by participants as feeling overwhelmed, exhausted and unable to continue living.⁵¹ Loss of control was also perceived to be caused by mobility, social status and social support losses.³² Once again, these losses led to feelings of helplessness where participants felt they no longer could continue living.^{32,51} The third qualitative study⁵⁰ identified deteriorating physical health and well-being and additional financial hardship as contributing to the self-harm episode or suicidal attempt. Despair and feelings of helplessness were also reported among participants that had attempted to end their lives.⁵⁰

Older adults mentioned increased feelings of loneliness and isolation, and these were major themes reported in the three qualitative studies.^{32,50,51} The previously described feelings of loss often resulted in participants feeling lonely and isolated.^{32,51} Participants also described having increased feelings of loneliness and isolation after the self-harm event, where family members regarded the episode as shameful.⁵⁰

Participants described and perceived ageing as burdensome, affecting all areas of daily living.^{32,50} Growing older was deemed to be a struggle and described with negative stereotypes of age and overall ageist views by older adults.³² Regret and missed opportunities were also voiced by participants as intensifying the felt internal struggle which contributed to the suicidal attempt.³²

(a)

	Study question	Population	Participation rate	Inclusion criteria	Sample size	Time frame	Blinded assessors	Repeated exposure	Defined outcomes	Loss to follow-up	Confounding
Armond <i>et al</i> 2017 ⁴⁷	✓	✓	-	✓	-	-	?	-	✓	-	X
Briskman <i>et al</i> 2017 ⁴⁴	✓	✓	-	✓	-	-	X	-	✓	-	X
Carter <i>et al</i> 2014 ³⁶	✓	✓	-	✓	-	-	?	-	✓	-	X
Cheung <i>et al</i> 2017 ³¹	✓	✓	?	✓	?	✓	?	✓	✓	✓	✓
Chiu <i>et al</i> 1996 ⁴⁰	✓	✓	✓	X	?	✓	X	✓	?	X	X
De Beer <i>et al</i> 2015 ²⁸	✓	✓	✓	✓	✓	✓	?	✓	✓	X	X
De Leo <i>et al</i> 2001 ⁵⁷	✓	✓	-	?	-	-	X	-	✓	-	X
De Leo <i>et al</i> 2002 ⁵⁸	✓	✓	✓	?	X	✓	X	✓	✓	X	X
Dennis <i>et al</i> 2007 ²⁰	✓	X	-	✓	-	-	?	-	✓	-	✓
Draper 1994 ³⁵	✓	✓	-	✓	-	-	X	-	X	-	X
Gavrielatos <i>et al</i> 2006 ⁴³	✓	✓	-	✓	-	-	X	-	✓	-	X
Gheshlaghi <i>et al</i> 2012 ⁴¹	✓	✓	-	✓	-	-	?	-	✓	-	X
Hawton <i>et al</i> 2006 ²¹	✓	✓	✓	✓	?	✓	✓	?	X	✓	✓
Hepple <i>et al</i> 1997 ³⁷	✓	✓	✓	✓	?	✓	X	✓	✓	X	X
Gokcelli <i>et al</i> 2017 ⁴⁸	✓	✓	-	✓	-	-	X	-	X	-	X
Kim <i>et al</i> 2011 ⁴²	✓	✓	-	✓	-	-	?	-	✓	-	X
Lawrence <i>et al</i> 2000 ³⁸	✓	✓	-	✓	-	-	?	-	✓	-	✓
Lamprecht <i>et al</i> 2005 ²²	✓	✓	✓	✓	?	✓	?	✓	✓	✓	✓
Lebret <i>et al</i> 2006 ⁴⁹	✓	✓	✓	✓	?	✓	X	✓	✓	✓	✓
Liu <i>et al</i> 2009 ⁵⁵	✓	✓	✓	✓	X	-	?	-	✓	-	✓
Logan <i>et al</i> 2007 ²⁹	✓	✓	-	✓	-	-	?	-	✓	-	X
Murphy <i>et al</i> 2012 ²³	✓	✓	✓	✓	?	✓	?	✓	✓	✓	✓
Nowers 1993 ²⁴	✓	✓	✓	✓	?	✓	?	✓	✓	✓	✓
Packer <i>et al</i> 2012 ¹⁹	✓	✓	-	✓	-	-	?	-	✓	-	X
Pierce 1987 ²⁵	✓	✓	✓	✓	?	✓	X	✓	✓	X	X
Pierce 1996 ²⁶	✓	✓	✓	✓	?	✓	?	✓	✓	X	X
Pillans <i>et al</i> 2017 ³⁹	✓	✓	-	✓	-	-	?	-	✓	-	✓
Ruths <i>et al</i> 2005 ²⁷	✓	✓	✓	✓	?	✓	?	✓	✓	X	X
Shah 2009 ³⁰	✓	✓	-	✓	-	-	?	-	✓	-	X
Takahashi <i>et al</i> 1995 ⁴⁵	✓	✓	-	✓	-	-	X	-	X	-	X
Ticehurst <i>et al</i> 2002 ³³	✓	✓	-	✓	-	-	?	-	X	-	✓
Tsoh <i>et al</i> 2005 ⁵³	✓	✓	X	✓	X	-	?	-	✓	-	✓
Van Orden <i>et al</i> 2015 ⁵²	✓	✓	✓	✓	-	✓	?	✓	✓	✓	✓
Wiktorsson <i>et al</i> 2010 ⁴⁶	✓	✓	-	✓	-	-	?	-	X	-	✓
Wynne <i>et al</i> 1987 ³⁴	✓	✓	-	✓	-	-	?	-	✓	-	X
Yang <i>et al</i> 2001 ⁵⁴	✓	✓	?	✓	-	-	X	-	✓	-	X
Zhang <i>et al</i> 2016 ⁵⁶	✓	✓	✓	✓	-	-	✓	-	✓	-	✓

(b)

	Study question	Methodology	Research design	Recruitment	Data collection	Relationship researcher	Ethical considerations	Data analysis	Findings	Research value
Bonnewyn <i>et al</i> 2014 ⁵¹	✓	✓	✓	✓	✓	?	✓	✓	✓	?
Crocker <i>et al</i> 2006 ³²	✓	✓	✓	✓	✓	?	✓	✓	✓	✓
Kim 2014 ⁵⁰	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

✓ Reported: element reported appropriately in study
 x Not reported: element not mentioned in study
 ? Cannot determine: lack of clarity to assess if element was reported
 - Not applicable: due to study design, element not applicable to report

Fig. 3 (a) Methodological quality assessment within studies. (b) Overall quality assessment across studies.




Fig. 3 (Continued)

Finally, participants also described feeling ‘too old’, leading them to their suicidal attempt to end the perceived ‘pain of old age’.⁵⁰

Influencing factors for self-harm in older adults

A thematic analysis of the influencing factors for self-harm in older adults (from the data presented in Table 1 from quantitative and qualitative studies) is summarised in Fig. 1. Influencing factors range from internal (e.g. age, gender) to external factors (e.g. financial worries, low education), showing the complex relationship between these factors throughout the presented layers. Figure 1 highlights the potential risk for self-harm and shows that it is not one single factor that independently influences self-harm in older adults. The themes are interconnected and layered across different individual, societal and healthcare settings.

Summary of findings

Overall, based on moderate quality evidence, previous history of self-harm, previous and current psychiatric treatment, and socio-demographic factors (single, living alone and younger older adults aged 60–74 years old) were found to be significant risk factors for self-harm repetition (Table 2). Others, such as alcohol/drug use, female gender, psychiatric history and a diagnosis of musculoskeletal conditions such as arthritis were also associated with self-harm repetition but the overall quality of evidence for these factors ranged from low to very low.

Discussion

This review presents current evidence regarding the characteristics of self-harm in older adults. Findings from this systematic review highlight self-harm in later life as having distinct characteristics to younger populations that should be explored to improve management and care for this age group. Despite sharing some characteristics of self-harm with younger populations (e.g. higher percentage in women, those with psychiatric history and those with a previous episode(s) of self-harm),^{12,59} there is an increased risk of repetition and suicide in older adults. Previous history of self-harm; previous and current psychiatric treatment; and sociodemographic factors including being single, living alone and being a younger older adult (60–74 years old) were more strongly associated with self-harm repetition.

Ranging from 19²⁹ to 65²³ yearly self-harm episodes per 100 000 people, findings from this review suggest prevalence rates to be lower compared with those reported in the literature of younger populations.^{60,61} However, the identified prevalence rates are to be taken with caution given that they are based on only seven studies which reported such findings, representing less than 5% of the total population of the systematic review. Furthermore, three of these studies^{25,27,28} have sample sizes of less than 200 participants, meaning their estimated rates must be taken with caution when calculating yearly self-harm rates per 100 000 people. There were also variant rates among the studies with only one study²⁹ identifying yearly rates of less than 20 per 100 000 people, with the rest of the studies having nearly double the number of rates. We believe the variance in rates could be attributed to the study design setting²⁹ and different healthcare system which reported non-suicidal self-injury as opposed to other presentations of self-harm (e.g. attempted suicide) as reported in the other studies. Furthermore, even with variant and lower prevalence rates compared with younger populations, the impact these presentations have on individuals and health services are significant. Time spent in hospital is longer in older adults who self-harm and medical complications are more likely, resulting in increased resource expenditure.^{29,62} Additionally, accuracy of self-harm estimates may not be completely representative given that the majority of the studies were based in hospital settings, and do not consider other presentations of self-harm which may not result in hospital attendance. With an increasing ageing population, it is important to acknowledge this possible under-representation of self-harm presentations in older adults. Older adults who self-harm are at a 67 times higher risk of suicide compared with younger populations.²³ This is congruent with worldwide epidemiological literature^{63,64} which states suicide rates in later life are one of the highest globally.

The use of self-poisoning is distinctive compared with other populations. One reason for this may be increased access to medication due to comorbid conditions that require prescribed medications. Nearly one third of the older adults were being prescribed antidepressants, giving them increased access to tablets for use in overdose. Data from the UK’s Office for National Statistics highlight that over one third of self-poisoning deaths were due to antidepressant overdose in 2014.^{65,66}

Findings suggest that older adults who self-harm report feelings of isolation, loneliness and loss of control. Ageing and reaching later life were perceived as burdensome by older adults, which

Table 3 Summary of findings on risk factors for self-harm repetition in older adults

		Evidence base	Strength association	Strength of evidence (GRADE) ^{a,b}	Comments
Sociodemographic factors	Female gender	Three studies ^{21,35,49} <i>n</i> = 858	Significance value $P < 0.05$ using χ^2 estimates $P = 0.014$, ²¹ 0.01 ⁴⁹	⊕⊕ Low	Uncertainty due to incomplete estimates presented in two of the studies
	Single, living alone and younger age (60–74 years old)	One study ²³ <i>n</i> = 1177	$P < 0.05$ (a) Single: hazard ratio 1.5, 95% CI 1.0–2.1 (b) Living alone: hazard ratio 1.5, 95% CI 1.0–2.3 (c) Younger age (60–74 years old): multivariate hazard ratio 1.8, 95% CI 1.2–2.8	⊕⊕⊕ Moderate	Evidence limited to one study, but strong association provided and large sample size
	No caregiver	One study ⁵⁶ <i>n</i> = 63	Odds ratio 1.82, 95% CI 1.04–3.33	⊕⊕ Low	Strong association provided but limited evidence with one study and small sample size
Clinical factors	Self-harm history	Two studies ^{23,53} <i>n</i> = 1240	Significance value $P < 0.05$ Multivariate hazard ratio ²³ 1.9, 95% CI 1.4–2.8 Adjusted odds ratio ⁵³ 32.9, 95% CI 3.2–339.37	⊕⊕⊕ Moderate	Mixed evidence for strength of association among studies. Stronger association found in study with increased number of participants
	Psychiatric history	Two studies ^{35,37} <i>n</i> = 169	$\chi^2 = 5.61$; $P < 0.05$ ³⁷	⊕ Very low	Uncertainty due to incomplete estimates presented
	Depression diagnosis	Four studies ^{37,53,55,56} <i>n</i> = 272	Adjusted odds ratio ⁵³ 59.2, 95% CI 6.4–546.6 Odds ratio ⁵⁵ 8.38, 95% CI 2.27–30.93 Odds ratio ⁵⁶ 5.19, 95% CI 2.92–9.22 $\chi^2 = 4.98$; $P < 0.05$ ³⁷	⊕⊕ Low	Despite multiple studies included, small sample size. Mixed evidence regarding strength of association, particularly imprecision of overall estimates
	Previous and current psychiatric treatment	Three studies ^{23,31,37} <i>n</i> = 1616	$P < 0.05$ Multivariate hazard ratio ²³ 1.8, 95% CI 1.2–2.7 Odds ratio ³¹ 2.73, 95% CI 1.20–6.25 $\chi^2 = 4.59$ ³⁷	⊕⊕⊕ Moderate	Strong association with estimates provided. Large sample size and multiple studies included
Other factors	Arthritis diagnosis	One study ⁵³ <i>n</i> = 66	Adjusted odds ratio 22.6, 95% CI 3.2–157.3	⊕ Very low	Low association provided with large imprecision in estimates. Limited evidence from only one study
	Time (12 months)	One study ²¹ <i>n</i> = 730	$P = 0.042$	⊕⊕ Low	Limited evidence from one study but large sample size
	Alcohol and drug use	Two studies ^{23,31} <i>n</i> = 1516	$P < 0.05$ Hazard ratio ²³ 1.9, 95% CI 1.5–5.1 Odds ratio ³¹ 3.87, 95% CI 1.35–11.12	⊕⊕ Low	Large sample size but inconsistency due to mixed results in strength of association among studies
	Poorer function of self-care ^c	Two studies ^{53,56} <i>n</i> = 129	Adjusted odds ratio ⁵³ 0.3, 95% CI 0.1–0.7 Odds ratio ⁵⁶ 0.83, 95% CI 0.76–0.92	⊕⊕ Low	Limited evidence with small sample size. Validity of tool used unknown

a. Modified Grading of Recommendations Assessment, Development and Evaluation (GRADE) system used to assess overall quality of risk factors. Elements used to assess evidence: risk of bias, inconsistency, indirectness, imprecision, large effect (strength of association) and dose-response gradient.

b. Meanings of symbols for quality of evidence across studies: ⊕⊕⊕⊕ High, further research is very unlikely to change our confidence in the estimate of effect; ⊕⊕⊕ Moderate, further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate; ⊕⊕ Low, further research is likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate; ⊕ Very low, any estimate of effect is very uncertain.

c. Measured using the Lawton Instrumental Activities of Daily Living Scale.

Table 4 Major themes with quotes from qualitative studies

Major themes	Bonnewyn <i>et al</i> , 2014 ⁵¹	Crocker <i>et al</i> , 2006 ³²	Kim, 2014 ⁵⁰
Loss of control leading to suicide attempt	'I was very tired. Completely exhausted. I did not see a way out anymore. I was tired doing the dishes, I was tired making up my bed. Tired, tired, always tired. But then again, I had almost not slept for months and months on end. I never slept during night time. My eyes hurt so much, I could simply not close them anymore.' (Female participant 1)	'Once I retired, I had no further aim, and had nothing to get up for. I didn't know what to do with my day. [...] My partner said "Just take that day as it comes, read the paper, go out and get the paper, do the chores ..." and I thought "Oh god, all unconstructive things".' (Male participant 1)	'I've tried twice to kill myself. It is not as easy to commit suicide as people think. I know it is a sin to do it, but I can't change my mind about ending this painful life. It's always stuck in my mind. I eat a lot of sugary foods and I do not take insulin because I think I have lived long enough.' (Female participant 5)
Increased loneliness and isolation	'Six months after my husband passed away, I really started to realize that ... I am on my own now. The children, they came in and I wanted ... I was troubled by something, I wanted to talk about it, but I could not.' (Female participant 2)	'When it got to the second stage [before attempt] it felt like that again you know, vanishing, you know and I thought I can't go on, on my own. And it's funny that because I've always been a loner.' (Female participant 3)	'I opened my eyes after three days. I was lying down in my room. No one knew what I had done. That was really sad and embarrassing. I felt terrible because no one cared about me.' (Female participant 6)
Ageing perceived as burdensome and affecting daily living	'It felt as if I could no longer cope. My greatest fear and biggest problem was that I would no longer be able to do the things which I was able to do before: looking after the grandchildren, washing, ironing, everything related to housekeeping. I am no longer able to do that and that is my biggest fear: that I won't be able to do that in the future. I can't do anything anymore, nothing works out; I am no longer of use. I can no longer do the things which I used to do before anyway.' (Female participant 1)	'Oh yes, I've been independent since I was born, let's put it that way. I never really depended on anybody or relied on anybody. I was very, very independent. I was a very feminine person, very sexy.' (Female participant 4)	'They [doctors] have been telling me I need surgery since last year. But why should I? To live longer? I don't have the money anyway. It would have been great if I had just died. This is more painful.' (Male participant 2)

contributed to their self-harm episode. However, these experiences were limited to the context of self-harm with exclusive suicidal intent.

Considerations for interpretation of findings

There are three main factors to consider when interpreting findings from this review. First, different terminologies were used across studies to refer to acts of self-harm, reflecting the ongoing heterogeneity of meanings inherent in the concept. For instance, definitions of self-harm in the literature included non-suicidal self-injury, self-harm and attempted suicide. Most of the included studies ($n = 29$) classified self-harm as attempted suicide, i.e. as holding an exclusively suicidal intent, which is not always the case.

Second, the design and reporting of many of the included studies did not allow for a comprehensive capture and statistical synthesis of all predefined outcomes (e.g. risk factors for repeated self-harm) as set out by the review. For instance, over half of the included studies were descriptive observational studies (e.g. cross-sectional) which mainly report disease distribution among populations to see whether a disease or condition is present or not.⁶⁷ This means that factors such as potential confounders and direction of causality between exposure and outcome could not always be determined for the whole older adult population. However, the availability of analytic study designs ($n = 14$ cohort studies) allowed more detailed exploration of the factors that influence self-harm in older adults. This is a strength for the evidence presented in this review as the inclusion of varied study designs ensured no evidence was lost and all available evidence is used to inform future research and practice.

Third, findings from this review are limited to data presented from included studies, which were predominantly based on self-harm presentations to hospital settings ($n = 34$). For instance, the yearly self-harm rates presented in this review were mostly based on studies conducted in hospital settings, as opposed to population- or community-based data. Not all self-harm episodes result in

hospital presentations, therefore other self-harm episodes (e.g. in the community) may not have been comprehensively captured in this review. Therefore, appropriate consideration must be taken when interpreting results from this review to avoid generalising to the wider population of older adults who self-harm.

Strengths and limitations of this review

This is the first review to systematically synthesise and appraise information regarding self-harm in older adults from both quantitative and qualitative studies. We believe reporting qualitative findings is of great importance to researchers and clinicians in the field, offering further explanation of self-harm in older adults. A further strength of this review is its emphasis on the inclusion of PPIE perspectives at all stages. An example of PPIE's collaboration in the review is the contribution to the development of Fig. 1, which was achieved by discussing the identified stressors with the PPIE group. As the National Institute of Health Research national advisory group INVOLVE⁶⁸ states, this makes research more relevant and likely to be addressing the needs of patients.

The conclusions of this review should be viewed with caution due to two factors. First, the majority of included studies were similar with regard to study setting, reporting self-harm in hospital settings rather than in the community. In addition to study selection by two independent pairs of reviewers, our search strategy was both sensitive and comprehensive, minimising the chances that any study might have been missed. Easier access to hospital patient records in the older adult population compared with conducting community-based research may explain the limited number of community-based studies. Another reason for the majority of evidence being predominantly from hospital settings may be the high level of stigma attached to self-harm,⁶⁹ resulting in resistance to help-seeking and/or accessing primary care services. Given the different settings and other factors influencing recording of self-harm, findings from the review may not be generalisable to the whole

population of older adults that self-harm, but mostly limited to a population of older adults attending hospital settings.

Second, evidence presented in systematic reviews is dependent on the inherent methodological quality of included studies. Despite quality assessment of the studies across domains being mostly moderate and with low risk of bias, the assessments highlighted certain areas of high risk of bias, including confounding, blinding of assessors and loss to follow-up. The low-quality rating of these areas is important to take into consideration when analysing the overall literature on self-harm in older adults.

Comparison with previous literature

Our review offers an update from previous reviews^{10,11} and explores factors not covered in previous work, such as self-harm repetition and motivations for self-harm. In contrast to other studies,^{10,11} we examined findings specifically around older populations and included additional study designs, i.e. both quantitative and qualitative studies. Other conducted reviews⁷⁰ assessing qualitative evidence may not be directly comparable to the present review given their inclusion of both direct and indirect self-harm. We adhere to the NICE guidelines definition of self-harm and view direct self-harm as distinct to indirect self-harm. This review therefore focused on direct self-harm only.

Furthermore, in younger populations there is empirical evidence which provides an explanation for underestimation of self-harm presentations.⁷¹ According to the iceberg model,⁷¹ there are three layers of self-harm presentations, with only two of them being overt and on the tip/surface of the iceberg: fatal self-harm (i.e. suicides) and hospital or clinical presentations of self-harm. However, the last and largest layer of the iceberg model⁷¹ is self-harm presentations in the community, which are mostly hidden given the lack of visibility. Therefore, it is likely that findings from this review can be translated to an iceberg model in older populations, once again highlighting the hidden element of self-harm and most likely underestimation of self-harm as found in this review.


Implications for clinical practice

Our findings are in line with NICE guideline CG16⁷² suggesting that identifying and managing older adults, whose self-harm is different from that in younger populations, is vital due to the increased risk of repetition and suicide in this population. Clinicians have the potential to intervene and prevent self-harm in older adults as frequent contact with health services is reported. This includes potential opportunities to reduce self-harm repetition (with resource implications, particularly for post-episode treatment), suicide and premature death.⁷³ In particular, it is important that clinicians prescribing antidepressants (among other medications) are aware of the increased risk of self-harm in this population and ensure adequate follow-up is in place. The model developed through this review offers the potential to inform clinicians about the possible influencing factors for self-harm in older adults (Fig. 1). However, it should not be used alone as other existing factors may not have been captured in the model given the limited hospital-based context of the majority of studies included in this review.

Future research

Further work is needed to identify appropriate resources and clear referral pathways to enable clinicians to support older people who self-harm. Future research should focus on populations of older adults engaging in self-harm within community settings and primary care so that there is a more comprehensive capture and understanding of self-harm in older adults, especially given that

most people who self-harm will be managed in primary care. Research exploring the different motivations for self-harm (suicidal or non-suicidal) would aid in clarifying the heterogeneous terminology used to refer to self-harm and further understand experiences of self-harm in later life. Lastly, data reporting standards within the psychiatric literature will benefit from careful consideration. Because of inadequate reporting/incomplete data provision across included studies, this review was unable to pool together findings in a meta-analysis. The agreement and compliance to high-reporting standards should be a priority for researchers and journals within the mental health field.

M. Isabela Troya, MSc, PhD Student , Research Institute for Primary Care and Health Sciences, Keele University, UK; **Opeyemi Babatunde**, PhD, Research Associate, Research Institute for Primary Care and Health Sciences, Keele University, UK; **Kay Polidano**, MSc, PhD Student, Research Institute for Primary Care and Health Sciences, Keele University, UK; **Bernadette Bartiam**, PhD, Senior Research Fellow, Family Medicine and Primary Care, Lee Kong Chian School of Medicine, Nanyang Technical University Singapore; **Erin McCloskey**, MA, MSc, PhD Student, School of Nursing, Midwifery and Social Work, Canterbury Christ Church University, UK; **Lisa Dikomitis**, PhD, Senior Lecturer in Sociology and Health, Research Institute for Primary Care and Health Sciences, Keele University, UK; **Carolyn A. Chew-Graham**, BSc, MBChB, MD, FRCGP, Professor of General Practice Research, Research Institute for Primary Care and Health Sciences, Keele University and West Midlands Collaboration for Leadership in Applied Health Research and Care; and Honorary Professor of Primary Care Mental Health, Midlands Partnership Foundation Trust, UK

Correspondence: M. Isabela Troya, Research Institute for Primary Care and Health Sciences, Keele University, David Weatherall Building, Staffordshire, ST5 5BG, UK. Email: m.i.troya.bermeo@keele.ac.uk

First received 8 Jul 2018, final revision 18 Dec 2018, accepted 21 Dec 2018

Supplementary material

Supplementary material is available online at <https://doi.org/10.1192/bjp.2019.11>.

Funding

M.I.T. and K.P. are funded by a Keele University ACORN studentship. Additional funding has been granted to M.I.T. by Santander Bank. C.A.C.-G. is partly funded by West Midlands Collaboration for Leadership in Applied Health Research and Care. E.M. is funded by a Canterbury Christ Church University MPhil/PhD scholarship in Health and Wellbeing.

Acknowledgements

We thank PPIE group members for their contribution to this review. We thank Echo and Brighter Futures. Special thanks to Dr Nadia Corp for assistance in the development of the search strategy.

References

- 1 Sinclair JM, Gray A, Rivero-Arias O, Saunders KE, Hawton K. Healthcare and social services resource use and costs of self-harm patients. *Soc Psychiatry Psychiatr Epidemiol* 2011; **46**: 263–71.
- 2 Haagsma JA, Graetz N, Bolliger I, Naghavi M, Higashi H, Mullany EC, et al. The global burden of injury: incidence, mortality, disability-adjusted life years and time trends from the Global Burden of Disease study 2013. *Inj Prev* 2016; **22**: 3–18.
- 3 Tsiachristas A, McDaid D, Casey D, Brand F, Leal J, Park AL, et al. General hospital costs in England of medical and psychiatric care for patients who self-harm: a retrospective analysis. *Lancet Psychiatry* 2017; **4**: 759–67.
- 4 National Institute for Health and Clinical Excellence (NICE). *Self-Harm in over 8s: Long-Term Management*. Clinical Guideline [CG133]. NICE, 2011 (<http://www.nice.org.uk/guidance/cg133>).
- 5 Hawton K, Zahl D, Weatherall R. Suicide following deliberate self-harm: long-term follow-up of patients who presented to a general hospital. *Br J Psychiatry* 2003; **182**: 537–42.
- 6 Sakinofsky I. Repetition of suicidal behaviour. In *The International Handbook of Suicide and Attempted Suicide* (eds K Hawton, K van Heeringen): 385–404. Wiley-Blackwell, 2000.

- 7 Office for National Statistics (ONS). *Overview of the UK Population: February 2016*. ONS, 2016 (<https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/articles/overviewoftheukpopulation/february2016/pdf>).
- 8 World Health Organization (WHO). *Preventing Suicide: A Global Imperative*. WHO, 2014 (http://www.who.int/mental_health/suicide-prevention/world_report_2014/en/).
- 9 World Health Organization (WHO). *Mental Health and Older Adults*. WHO, 2016 (www.who.int/mediacentre/factsheets/fs381/en/).
- 10 Chan J, Draper B, Banerjee S. Deliberate self-harm in older adults: a review of the literature from 1995 to 2004. *Int J Geriatr Psychiatry* 2007; **22**: 720–32.
- 11 Draper B. Editorial Review. Attempted suicide in old age. *Int J Geriatr Psychiatry* 1996; **11**: 577–87.
- 12 Morgan C, Webb RT, Carr MJ, Kontopantelis E, Green J, Chew-Graham CA, et al. Incidence, clinical management, and mortality risk following self harm among children and adolescents: cohort study in primary care. *BMJ* 2017; **359**: 435–51.
- 13 Jinks C, Carter P, Rhodes C, Taylor R, Beech R, Dziedzic K, et al. Patient and public involvement in primary care research—an example of ensuring its sustainability. *Res Involv Engagem* 2016; **2**: 1.
- 14 Given LM (ed.). *The SAGE Encyclopaedia of Qualitative Research Methods*. SAGE Publications, 2008.
- 15 Popay J, Roberts H, Sowden A, Petticrew M, Arai L, Rodgers M, et al. *Guidance on the Conduct of Narrative Synthesis in Systematic Reviews: A product from the ESRC Methods Programme*. Version 1: p. 92. Lancaster University, 2006.
- 16 Thomas J, Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Med Res Methodol* 2008; **8**: 45.
- 17 National Institutes of Health (NIH). *Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies*. NIH, 2018 (<https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools>).
- 18 Critical Appraisal Skills Programme (CASP). *CASP Qualitative Checklist*. CASP, 2018 (<https://casp-uk.net/wp-content/uploads/2018/01/CASP-Qualitative-Checklist.pdf>).
- 19 Packer J, Hussain MA, Shah SH, Srinivasan JR. Deliberate self-harm and the elderly: a volatile combination—an overview from the plastic surgery perspective. *Plast Surg Int* 2012; **2012**: 1–4.
- 20 Dennis MS, Wakefield P, Molloy C, Andrews H, Friedman T. A study of self-harm in older people: mental disorder, social factors and motives. *Aging Ment Health* 2007; **11**: 520–5.
- 21 Hawton K, Harriss L. Deliberate self-harm in people aged 60 years and over: characteristics and outcome of a 20-year cohort. *Int J Geriatr Psychiatry* 2006; **21**: 572–81.
- 22 Lamprecht HC, Pakrasi S, Gash A, Swann AG. Deliberate self-harm in older people revisited. *Int J Geriatr Psychiatry* 2005; **20**: 1090–6.
- 23 Murphy E, Kapur N, Webb R, Purandare N, Hawton K, Bergen H, et al. Risk factors for repetition and suicide following self-harm in older adults: multi-centre cohort study. *Br J Psychiatry* 2012; **200**: 399–404.
- 24 Nowers M. Deliberate self-harm in the elderly: a survey of one London Borough. *Int J Geriatr Psychiatry* 1993; **8**: 609–14.
- 25 Pierce D. Deliberate self-harm in the elderly. *Int J Geriatr Psychiatry* 1987; **2**: 105–10.
- 26 Pierce D. Deliberate self-harm: how do patients view their treatment? *Br J Psychiatry* 1996; **149**: 624–6.
- 27 Ruths FA, Tobiansky RI, Blanchard M. Deliberate self-harm (DSH) among older people: a retrospective study in Barnet, North London. *Int J Geriatr Psychiatry* 2005; **20**: 106–12.
- 28 de Beer WA, Murtagh J, Cheung G. Late-life self-harm in the Waikato Region. *Ethnicity* 2015; **4**: 17–3.
- 29 Logan J, Crosby A, Ryan G. Nonfatal self-inflicted injuries among adults aged ≥65 years—United States, 2005. *MMWR Morb Mortal Wkly Rep* 2007; **56**: 989–93.
- 30 Shah A. Attempted suicide in the elderly in England: age-associated rates, time trends and methods. *Int Psychogeriatr* 2009; **21**: 889–95.
- 31 Cheung G, Foster G, de Beer W, Gee S, Hawkes T, Rimkeit S, et al. Predictors for repeat self-harm among older people. *Int Psychogeriatr* 2017; **29**: 1237–45.
- 32 Crocker L, Clare L, Evans K. Giving up or finding a solution? The experience of attempted suicide in later life. *Aging Men Health* 2006; **10**: 638–47.
- 33 Ticehurst S, Carter GL, Clover KA, Whyte IM, Raymond J, Fryer J. Elderly patients with deliberate self-poisoning treated in an Australian general hospital. *Int Psychogeriatr* 2002; **14**: 97–105.
- 34 Wynne H, Bateman DN, Hassanyeh F, Rawlins MD, Woodhouse KW. Age and self-poisoning: the epidemiology in Newcastle upon Tyne in the 1980s. *Hum Toxicol* 1987; **6**: 511–5.
- 35 Draper B. Suicidal behaviour in the elderly. *Int J Geriatr Psychiatry* 1994; **9**: 655–61.
- 36 Carter MW, Reymann MR. ED use by older adults attempting suicide. *Am J Emerg Med* 2014; **32**: 535–40.
- 37 Hepple J, Quinton C. One hundred cases of attempted suicide in the elderly. *Br J Psychiatry* 1997; **171**: 42–6.
- 38 Lawrence D, Almeida OP, Hulse GK, Jablensky AV, Holman CAJ. Suicide and attempted suicide among older adults in Western Australia. *Psychological Med* 2000; **30**: 813–21.
- 39 Pillans PI, Page CB, Ilango S, Kashchuk A, Isbister GK. Self-poisoning by older Australians: a cohort study. *Med J Aust* 2017; **206**: 164–9.
- 40 Chiu HF, Lam LC, Pang AH, Leung CM, Wong CK. Attempted suicide by Chinese elderly in Hong Kong. *Gen Hosp Psychiatry* 1996; **18**: 444–7.
- 41 Gheshlaghi F, Salehi MJ. Suicide attempts by self-poisoning in elderly. *J Res Med Sci* 2012; **2**: 272–5.
- 42 Kim YR, Choi KH, Oh Y, Lee HK, Kweon YS, Lee CT, et al. Elderly suicide attempters by self-poisoning in Korea. *Int Psychogeriatr* 2011; **23**: 979–85.
- 43 Gavrielatos G, Komitopoulos N, Kanellos P, Varsamis E, Kogeorgos J. Suicidal attempts by prescription drug overdose in the elderly: a study of 44 cases. *Neuropsychiatr Dis Treat* 2006; **2**: 359–63.
- 44 Briskman I, Shelef A, Berger U, Baruch Y, Bar G, Asherov J, et al. Deliberate self-harm in older adults: a general hospital emergency department survey. *Israel Med Assoc J* 2017; **19**: 160–3.
- 45 Takahashi Y, Hirasawa H, Koyama K, Asakawa O, Kido M, Onose H, et al. Suicide and aging in Japan: an examination of treated elderly suicide attempters. *Int Psychogeriatr* 1995; **7**: 239–51.
- 46 Wiktorsson S, Runeson B, Skoog I, Östling S, Waern M. Attempted suicide in the elderly: characteristics of suicide attempters 70 years and older and a general population comparison group. *Am J Geriatr Psychiatry* 2010; **18**: 57–67.
- 47 Armond JDE, Armond RDE, Pereira TC, Chinaia C, Vendramini TL, Rodrigues CL. Self-injury and suicide attempt among the elderly population in the city of São Paulo. *J Bras Psiquiatr* 2017; **66**: 83–8.
- 48 Gokceli DK, Tasar PT, Akcam NO, Sahin S, Akarca FK, Aktas EO, et al. Evaluation of attempted older adults suicides admitted to a University Hospital Emergency Department: Izmir study. *Asian J Psychiatr* 2017; **30**: 196–9.
- 49 Lebre S, Perret-Vaille E, Mulliez A, Gerbaud L, Jalenques I. Elderly suicide attempters: characteristics and outcome. *Int J Geriatr Psychiatry* 2006; **21**: 1052–9.
- 50 Kim Y. Understanding the life experiences of older adults in Korea following a suicide attempt. *Qual Health Res* 2014; **24**: 1391–9.
- 51 Bonnewyn A, Shah A, Bruffaerts R, Schoevaerts K, Rober P, Van Parys H, et al. Reflections of older adults on the process preceding their suicide attempt: a qualitative approach. *Death Stud* 2014; **38**: 612–8.
- 52 Van Orden KA, Wiktorsson S, Duberstein P, Berg AI, Fässberg MM, Waern M. Reasons for attempted suicide in later life. *Am J Geriatr Psychiatry* 2015; **23**: 536–44.
- 53 Tsoh J, Chiu HF, Duberstein PR, Chan SS, Chi I, Yip PS, et al. Attempted suicide in elderly Chinese persons: a multi-group, controlled study. *Am J Geriatr Psychiatry* 2005; **13**: 562–71.
- 54 Yang CH, Tsai SJ, Chang JW, Hwang JP. Characteristics of Chinese suicide attempters admitted to a geropsychiatric unit. *Int J Geriatr Psychiatry* 2001; **16**: 1033–6.
- 55 Liu IC, Chiu CH. Case-control study of suicide attempts in the elderly. *Int Psychogeriatr* 2009; **21**: 896–902.
- 56 Zhang W, Ding H, Su P, Xu Q, Du L, Xie C, et al. Prevalence and risk factors for attempted suicide in the elderly: a cross-sectional study in Shanghai, China. *Int Psychogeriatr* 2016; **29**: 709–15.
- 57 De Leo D, Padoani W, Scocco P, Lie D, Bille-Brahe U, Arensman E, et al. Attempted and completed suicide in older subjects: results from the WHO/EURO Multicentre Study of Suicidal Behaviour. *Int J Geriatr Psychiatry* 2001; **16**: 300–10.
- 58 De Leo D, Padoani W, Lonnqvist J, Kerkhof AJ, Bille-Brahe U, Michel K, et al. Repetition of suicidal behaviour in elderly Europeans: a prospective longitudinal study. *J Affect Disord* 2002; **72**: 291–5.
- 59 Hawton K, Rodham K, Evans E, Weatherall R. Deliberate self-harm in adolescents: self-report survey in schools in England. *BMJ* 2002; **325**: 1207–11.
- 60 Muehlenkamp JJ, Claes L, Havertape L, Plener PL. International prevalence of adolescent non-suicidal self-injury and deliberate self-harm. *Child Adolesc Psychiatry Ment Health* 2012; **6**: 10.
- 61 Swannell SV, Martin GE, Page A, Hasking P, St John NJ. Prevalence of nonsuicidal self-injury in nonclinical samples: systematic review, meta-analysis and meta-regression. *Suicide Life-Threat Behav* 2014; **44**: 273–303.

- 62 Czernin S, Vogel M, Flückiger M, Muheim F, Bourgnon JC, Reichelt M, et al. Cost of attempted suicide: a retrospective study of extent and associated factors. *Swiss Med Wkly* 2012; **142**: w13648.
- 63 Bertolote JM, Fleischmann A. A global perspective in the epidemiology of suicide. *Suicidologi* 2015; **7**: 6–8.
- 64 Shah A. The relationship between suicide rates and age: an analysis of multinational data from the World Health Organization. *Int Psychogeriatr* 2007; **19**: 1141–52.
- 65 Office for National Statistics (ONS) *Deaths Related to Drug Poisoning in England and Wales 2014–2015, 2015* (<https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/deathsrelatedtodrugpoisoninginenglandandwales/2015-09-03#antidepressants>).
- 66 Office for National Statistics (ONS) *Suicides in the United Kingdom: 2014 Registrations, 2016* (<https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/suicidesintheunitedkingdom/2014registrations#uk-suicides>).
- 67 Hennekens CH, Buring JE, Mayrent SL. *Epidemiology in medicine*. Little Brown and Company, 1987.
- 68 INVOLVE. *Public Involvement in Systematic Reviews: Supplement to the Briefing Notes for Researchers*. INVOLVE, 2012 (https://www.invo.org.uk/wp-content/uploads/2014/10/INVOLVEPublicInvolvementSystematicReviews_updated-July-2014_WEB.pdf).
- 69 McAllister M. Multiple meanings of self-harm: a critical review. *Int J Ment Health Nurs* 2003; **12**: 177–85.
- 70 Wand APF, Peisah C, Draper B, Brodaty H. Understanding self-harm in older people: a systematic review of qualitative studies. *Aging Ment Health* 2018; **22**: 289–98.
- 71 Geulayov G, Casey D, McDonald KC, Foster P, Pritchard K, Wells C, et al. Incidence of suicide, hospital-presenting non-fatal self-harm, and community-occurring non-fatal self-harm in adolescents in England (the iceberg model of self-harm): a retrospective study. *Lancet Psychiatry* 2018; **5**: 167–74.
- 72 National Institute for Clinical Excellence (NICE). *Self-Harm: The Short-Term Physical and Psychological Management and Secondary Prevention of Self-Harm in Primary and Secondary Care*. Clinical Guideline [CG16]. NICE, 2004 (<https://www.nice.org.uk/guidance/cg16>).
- 73 Carr MJ, Ashcroft DM, Kontopantelis E, While D, Awenat Y, Cooper J, et al. Premature death among primary care patients with a history of self-harm. *Ann Fam Med* 2017; **15**: 246–54.

