



# COVID-19 vaccination

# A guide to phase 2 of the programme

This leaflet explains phase 2 of the COVID-19 vaccination programme, who is eligible and who needs to have the vaccine to protect them from Coronavirus.

Find out more at [nhs/COVIDVaccine](https://nhs.uk/COVIDVaccine)

## What is COVID-19 or Coronavirus?

COVID-19 is caused by a new coronavirus, known as SARS-CoV-2. It was first identified in late 2019. It is very infectious and can lead to severe respiratory disease.

Many people who are infected may not have any symptoms or only have mild symptoms. These commonly start with cough, fever, headache and loss of taste or smell.

Some people will feel very tired, have aching muscles, diarrhoea and vomiting, fever and confusion. A small number of people then go on to have severe disease which may require hospitalisation or admission to intensive care.

**Overall fewer than 1 in 100 people who are infected will die from COVID-19, but in those over 75 years of age this rises to 1 in 10.**

There is no cure for COVID-19 although some newly tested treatments do help to reduce the risk of complications.

## About the types of vaccine

In the UK several different types of COVID-19 vaccines will be used during 2021. The vaccines will only be approved on the basis of large studies of safety and effectiveness.

Over 30 million people in the UK have been vaccinated against COVID-19 and the vaccine has already prevented many cases and more than 6 thousand deaths from COVID-19.

Recently there have been reports of an extremely rare condition involving blood clots and unusual bleeding after vaccination with AstraZeneca (AZ). This is being carefully reviewed but the risk factors for this condition are not yet clear. Because of the high risk of complications and death from COVID-19, the MHRA, the World Health Organisation and the European Medicines Agency have concluded that the balance is very much in favour of vaccination.

## Who should have the COVID-19 vaccines?

In the first phase of the programme the Joint Committee on Vaccination and Immunisation (JCVI), an independent expert group, recommended that the NHS offers these vaccines to those at highest risk of catching the infection and of suffering serious complications if they catch the infection.

This includes older adults, frontline health and social care workers, care home residents and staff, and those with certain clinical conditions.

## Am I at risk from COVID-19 infection?

Coronavirus can affect anyone. If you are an older adult and have a long-term health condition, you should have already been offered the vaccine.

## Who is now eligible for the vaccination?

The vaccine is now being offered to people at lower risk from the complications of COVID-19. You are still at risk of catching the infection and passing it on to others.

For most younger people COVID-19 is usually a milder illness that rarely leads to complications. For a few people the symptoms may last for longer than the usual two to three weeks. The vaccination will help to protect you against COVID-19.

In the second phase of the programme the vaccine will be offered to those under 50 years old in the following order:

- 40 - 49 years
- 30 - 39 years
- 18 years and over

The risk remains higher in older ages. For example the risk of dying from COVID-19 in someone aged 40 - 49 is 3 times higher than someone in the 30 to 39 year age group and 12 times higher than someone in the 20 to 29 year age group.





As the programme progresses, the JCVI will advise on the appropriate vaccine for each age group. The vaccine offered may depend on your age. Currently the JCVI has advised it is preferable for healthy people under 30 to have a vaccine other than AstraZeneca (AZ). The time when it will be offered will be based on the availability of those vaccines. Anybody who was previously eligible for vaccination remains so and should come forward to start or complete their recommended course.

### Who cannot have the vaccine?

The vaccines do not contain organisms that grow in the body, and so are safe for people with disorders of the immune system. These people may not respond so well to the vaccine. A very small number of people who are at risk of COVID-19 cannot have the vaccine – this includes some people who have severe allergies.

### Who should wait to have the vaccine?

If you are currently unwell, self-isolating, or waiting for a COVID-19 test you should delay vaccination until later. Pregnant women may prefer to wait to have the vaccine until they have completed their pregnancy. Read the detailed information on [www.nhs.uk/covidvaccination](http://www.nhs.uk/covidvaccination).

### Will the vaccine protect me?

The COVID-19 vaccination will reduce the chance of you suffering from COVID-19 disease. It may take a few weeks for your body to build up some protection from the vaccine. Like all medicines, no vaccine is completely effective – some people may still get COVID-19 despite having a vaccination, but this should be less severe.

**If you need more information on symptoms visit**  
[www.nhs.uk/conditions/coronavirus-covid-19/symptoms](http://www.nhs.uk/conditions/coronavirus-covid-19/symptoms)

## Will the vaccine have side effects?

Like all medicines, vaccines can cause side effects. Most of these are mild and short-term, and not everyone gets them. Even if you do have symptoms after the first dose, you still need to have the second dose. Although you should get good protection from the first dose, having the second dose should give you longer lasting protection against the virus.

Very common side effects include:

- having a painful, heavy feeling and tenderness in the arm where you had your injection. This tends to be worst around 1-2 days after the vaccine
- feeling tired
- headache
- general aches, or mild flu like symptoms

You can rest and take the normal dose of paracetamol (follow the advice in the packaging) to help you feel better. Although feeling feverish is not uncommon for two to three days, a high temperature is unusual and may indicate you have COVID-19 or another infection. Symptoms following vaccination normally last less than a week. If your symptoms seem to get worse or if you are concerned, call NHS 111.

### **If you experience any of the following from around 4 days to four weeks after vaccination you should seek medical advice urgently.**

- a new, severe headache which is not helped by usual painkillers or is getting worse
- an unusual headache which seems worse when lying down or bending over or may be accompanied by
  - blurred vision, nausea and vomiting
  - difficulty with your speech,
  - weakness, drowsiness or seizures
- new, unexplained pinprick bruising or bleeding
- shortness of breath, chest pain, leg swelling or persistent abdominal pain



If you have any of the above symptoms after your first vaccination, you should speak to your doctor or specialist before having the second dose.

If you do seek advice from a doctor or nurse please make sure you tell them about your vaccination (show them the vaccination card if possible) so that they can assess you properly. You can also report suspected side effects to vaccines and medicines through the Yellow Card scheme. See page 8.

**If you are currently in the clinically extremely vulnerable group, please continue to follow the government guidance.**

### What can I do after I have had the COVID-19 vaccine?

The vaccine cannot give you COVID-19 infection, and two doses will reduce your chance of becoming seriously ill. We do not yet know how much it will reduce the chance of you catching and passing on the virus. So, it is important to continue to protect those around you.

Remember to protect yourself and your family, friends and colleagues you MUST still follow any national or local restrictions and:

- practise social distancing
- wear a face mask
- wash your hands regularly
- open windows to let fresh air in
- follow the current guidance at [www.gov.uk/coronavirus](http://www.gov.uk/coronavirus)



## Can I catch COVID-19 from the vaccine?

You cannot catch COVID-19 from the vaccine but it is possible to have caught COVID-19 and not realise you have the symptoms until after your vaccination appointment.

The most important symptoms of COVID-19 are recent onset of any of the following:

- a new continuous cough
- a high temperature
- a loss of, or change in your normal sense of taste or smell

If you have the symptoms above, stay at home and arrange to have a test.

## What do I do next?

After you have had the first dose you need to plan to attend your second appointment. You should have a record card and your next appointment should be between 3 and 12 weeks later.

Although the first dose will give you good protection, you need the second dose to get longer lasting protection.

## What should I do if I am not well when it is my next appointment?

If you are unwell, it is better to wait until you have recovered to have your vaccine, but you should try to have it as soon as possible.

You should not attend a vaccine appointment if you are self-isolating, waiting for a COVID-19 test or within 4 weeks of having a positive COVID-19 test.

**Keep your card safe  
and make sure you keep  
your next appointment  
to get your second dose.**

Don't forget your  
COVID-19 vaccination **NHS**

**Make sure you keep this record  
card in your purse or wallet**

For more information on the  
COVID-19 vaccination or what  
to do after your vaccination,  
see [www.nhs.uk/covidvaccine](http://www.nhs.uk/covidvaccine)

**COVID-19 immunisation**  
Protect yourself.



Please read the product information leaflet for more details on your vaccine, including possible side effects, by searching Coronavirus Yellow Card. You can also report suspected side effects on the same website or by downloading the Yellow Card app.



[coronavirus-yellowcard.mhra.gov.uk](https://coronavirus-yellowcard.mhra.gov.uk)

## Remember

COVID-19 is spread through droplets expelled from the nose or mouth, particularly when speaking or coughing. It can also be picked up by touching your eyes, nose and mouth after contact with contaminated objects and surfaces.

You **MUST** still continue to follow any national or local restrictions and:

- practise social distancing
- wear a face mask
- wash your hands regularly
- open windows to let fresh air in
- follow the current guidance at [www.gov.uk/coronavirus](https://www.gov.uk/coronavirus)

## Vaccination, helping to protect those most vulnerable.

If you need more information on the COVID-19 vaccination please visit: [nhs.uk/CoronavirusVaccination](https://nhs.uk/CoronavirusVaccination)

You can order more of these free leaflets at [www.healthpublications.gov.uk](https://www.healthpublications.gov.uk)

© Crown copyright 2021. COV2020351V4EN 10M 2p MAR (APS) 2021  
Public Health England gateway number: 2020616